Theory for long range magnetic order in nanometer films

Cong B.T., Thao P.H., Cong P.T.

Faculty of Physics, Hanoi University of Science, 334 Nguyen Trai, Hanoi, Viet Nam; Physikalisches Institut, Universität Frankfurt, SFB/TR49, Frankfurt(M), Germany

Abstract: The functional integral method and Heisenberg model for localised spins were used for the calculation of the thermodynamics characteristics of magnetic films. It was shown in the Gaussian approximation that the Curie temperature is strongly reduced when the film's thickness reduces to the nanometer region. © 2010 IOP Publishing Ltd.

Authors with affiliations:

• Cong, B.T., Faculty of Physics, Hanoi University of Science, 334 Nguyen Trai, Hanoi, Viet Nam

• Thao, P.H., Faculty of Physics, Hanoi University of Science, 334 Nguyen Trai, Hanoi, Viet Nam

• Cong, P.T., Faculty of Physics, Hanoi University of Science, 334 Nguyen Trai, Hanoi, Viet Nam, Physikalisches Institut, Universität Frankfurt, SFB/TR49, Frankfurt(M), Germany

References:

• Gajdzik, M., Trappman, T., Sürgers, C., Löhneysen, H.V., (1998) Phys. Rev., 57 (6), p. 3525

• Huang, F., Kief, M.T., Mankey, G.J., Willis, R.F., (1994) Phys. Rev., 49 (6), p. 3962

• Tjeng, L.H., Izderda, Y.U., Rudoff, P., Sette, F., Chen, C.T., (1992) J. Magn. Magn. Mater., 109 (2-3), p. 288

• Fischer, M.E., Barber, M.N., (1972) Phys. Rev. Lett., 28 (23), p. 1516

• Lang, X.Y., Zheng, W.T., Jiang, Q., (2006) Phys. Rev., 73 (22), p. 224444

• Sun, C.Q., Zhong, W.H., Li, S., Tay, B.K., Bai, H.I., Jiang, E.Y., (2004) J. Phys. Chem., 108 (3), p. 1080

• Varkarchuk, I.A., Rudavskii, Y.K., (1981) Teor. Mat. Fiz., 49, p. 235. , in Russian

• Thanh Cong, B., (1986) Phys. Status Solidi, 134 (2), p. 569

• Zaidi, M.H., (1983) Fortschr. Phys., 31 (7), p. 403

• Izuymov, Y.A., Kassanogly, F.A., Skriabin, Y.N., (1974) Field Methods in the Theory of Ferromagnets, , Mir: Moscow