Chemical composition of the essential oil from the rhizomes of Alpinia henryi K. Schum. of Vietnam

Phan G.M., Phan S.T., Konig W.A.

Faculty of Chemistry, College of Natural Science, Vietnam National University, 19 Le Thanh Tong Street, Hanoi, Viet Nam; Institut für Organische Chemie, Universität Hamburg, D-20146 Hamburg, Germany

Abstract: The essential oil from the fresh rhizomes of Alpinia henryi K. Schum. (Zingiberaceae) collected in Vietnam was analyzed by GC and GC/MS. Twenty-five compounds, representing 86.4% of the oil, were identified. The oil was characterized by a high content of 1,8-cineole (45.1%). © 2007 Allured Publishing Corp.

Author Keywords: 1,8-cineole; Alpinia henryi; Essential oil composition; Zingiberaceae Index Keywords: Hydrocarbons; Plants (botany); 1,8-cineole; Alpinia henryi; Zingiberaceae; Essential oils; Alpinia; Zingiberaceae

Year: 2007

Source title: Journal of Essential Oil Research

Volume: 19 Issue: 6

Page: 507-508 Link: Scorpus Link

Correspondence Address: Phan, S.T.; Faculty of Chemistry, College of Natural Science, Vietnam National

University, 19 Le Thanh Tong Street, Hanoi, Viet Nam

ISSN: 10412905 CODEN: JEORE

Language of Original Document: English

Abbreviated Source Title: Journal of Essential Oil Research

Document Type: Article

Source: Scopus

Authors with affiliations:

- Phan, G.M., Faculty of Chemistry, College of Natural Science, Vietnam National University, 19 Le Thanh Tong Street, Hanoi, Viet Nam
- Phan, S.T., Faculty of Chemistry, College of Natural Science, Vietnam National University, 19 Le Thanh Tong Street, Hanoi, Viet Nam
- König, W.A., Institut für Organische Chemie, Universität Hamburg, D-20146 Hamburg, Germany

References:

• D.H. Hochmuth, W.A. König and D. Joulain, MassFinder 2.3. Software & Data Bank, Hamburg (2003). Available at: www.massfinder.com