## Distribution of persistent organic pollutants and polycyclic aromatic hydrocarbons in sediment samples from Vietnam

Kishida M., Imamura K., Maeda Y., Lan T.T.N., Thao N.T.P., Pham H.V.

Environmental Pollution Control Center, Osaka Prefectural Government, 1-3-62 Nakamichi, Higashinari-ku, Osaka 537-0025, Japan; Graduate School of Engineering, Osaka Prefecture University, 1-1 Gakuen-cho, Naka-ku, Sakai, Osaka 599-8531, Japan; College of Natural Science, Vietnam National University of Ho Chi Minh City, 227 Nguyen Van Cu Str., Dist.5, Ho Chi Minh City, Viet Nam; National Center for Natural Science and Technology, Hoang Quoc Viet Str., Hghia Do, Cau Giay, Hanoi, Viet Nam; College of Science, Vietnam National University of Hanoi, T3 Building, 333 N. Trai Str., Thanh Xuan Dist., Hanoi, Viet Nam

Abstract: The presence of eight kinds of persistent organic pollutants (POPs) such as DDT and its metabolites (DDTs), hexachlorocyclohexanes (HCHs), chlordane compounds (CHLs), drin compounds (Drins), heptachlor, hexachlorobenzene (HCB), heptachlor-epoxide, polychlorinated biphenyls (PCBs) and sixty-four polycyclic aromatic hydrocarbon compounds (PAHs) was identified using high resolution gas chromatography/high resolution mass spectrometry (HRGC/HRMS) to investigate their distribution in surface sediment from Hanoi, Hue, and Ho Chi Minh in Vietnam. A survey of sediment samples from Osaka was conducted for comparison. The concentrations of  $\Sigma$ DDTs,  $\Sigma$ CHLs,  $\Sigma$ PCBs and  $\Sigma$ PAHs in Vietnam were 0.19-140, N.D.-9.0, 0.11-110, and 30-5500 ng/g-dry, respectively. Concentrations of these compounds in urban areas were higher than those in other areas. In addition, the  $\Sigma$ DDT concentrations in Vietnamese urban areas were higher than those in Osaka. These results suggest that most DDTs would be used as insecticides for the purpose of health services rather than as agricultural chemicals. PAH pollution in urban areas and suburbs is caused mainly by runoff of petrol, whereas in rural areas, the combustion of fossil fuels and biomass is the major pollutant source.

Author Keywords: High resolution gas chromatography/high resolution mass spectrometry; Persistent organic pollutant; Polycyclic aromatic hydrocarbon; Sediment sample; Vietnam

Index Keywords: fossil fuel; polycyclic aromatic hydrocarbon; article; biomass; gas chromatography; health service; mass spectrometry; organic pollution; urban area; Viet Nam

Year: 2007 Source title: Journal of Health Science Volume: 53 Issue: 3 Page : 291-301 Cited by: 5 Link: Scorpus Link Correspondence Address: Kishida, M.; Environmental Pollution Control Center, Osaka Prefectural Government, 1-3-62 Nakamichi, Higashinari-ku, Osaka 537-0025, Japan; email: KishidaMas@mbox.pref.osaka.lg.jp ISSN: 13449702 CODEN: JHSCF

DOI: 10.1248/jhs.53.291

Language of Original Document: English

Abbreviated Source Title: Journal of Health Science

Document Type: Article

Source: Scopus

Authors with affiliations:

- Kishida, M., Environmental Pollution Control Center, Osaka Prefectural Government, 1-3-62 Nakamichi, Higashinari-ku, Osaka 537-0025, Japan
- Imamura, K., Environmental Pollution Control Center, Osaka Prefectural Government, 1-3-62 Nakamichi, Higashinari-ku, Osaka 537-0025, Japan
- Maeda, Y., Graduate School of Engineering, Osaka Prefecture University, 1-1 Gakuen-cho, Naka-ku, Sakai, Osaka 599-8531, Japan
- Lan, T.T.N., College of Natural Science, Vietnam National University of Ho Chi Minh City, 227 Nguyen Van Cu Str., Dist.5, Ho Chi Minh City, Viet Nam
- Thao, N.T.P., National Center for Natural Science and Technology, Hoang Quoc Viet Str., Hghia Do, Cau Giay, Hanoi, Viet Nam
- Pham, H.V., College of Science, Vietnam National University of Hanoi, T3 Building, 333 N. Trai Str., Thanh Xuan Dist., Hanoi, Viet Nam

References:

- Halsall, C.J., Bailey, R., Stern, G.A., Barrie, L.A., Fellin, P., Muir, D.C.G., Rosenberg, B., Pastukhov, B., Multi-year observations of organohalogen pesticides in the Arctic atmosphere (1998) Environ. Pollut, 102, pp. 51-62
- Iwata, H., Tanabe, S., Sakai, N., Nishimura, A., Tatsukawa, R., Geographical distribution of persistent organochlorine in air, water, and sediments from Asia and Oceania, and their implication for global distribution from lower latitudes (1994) Environ. Pollut, 85, pp. 15-33
- Iwata, H., Tanabe, S., Sakai, N., Tatsukawa, R., Distribution of persistent organochlorines in the oceanic air and surface seawater and the role of ocean on their global transport and fate (1993) Environ. Sci. Technol, 27, pp. 1080-1098
- Minh, T. B., Minh, N. H., Kunisue, T., Watanabe, M., Iwata, H., Pham, H. V., Hue, D. H., Qui, V., Tuyen, B. C. and Tanabe, S. (2003) Persistent organic pollutants (POPs) in Vietnamese environment A review of contamination, fate, and toxic potential. In Proceedings of environmental science and technology for sustainable development, the 4th General Seminar of the Core University Program, July 14-15, Osaka, Japan, pp.63-71Nhan, D.D., Am, N.M., Carvalho, F.P., Villeneuve, J.-P., Cattini, C., Organochlorine pesticides and PCBs along the coast of North Vietnam (1999) Sci. Total Environ, 237 -8, pp. 363-371
- Nhan, D.D., Carvalho, F.P., Am, N.M., Tuan, N.Q., Yen, N.T.H., Villeneuve, J.-P., Cattini, C., Chlorinated pesticides and PCBs in sediments and mollusks from freshwater canals in the Hanoi region (2001) Environ. Pollut, 112, pp. 311-320
- Kannan, K., Tanabe, S., Tatsukawa, R., Geographical distribution and accumulation features of organochlorine residues in fish from tropical Asia and Oceania (1995) Environ. Sci. Technol, 29, pp. 2673-2683
- Zakaria, M.P., Takada, H., Tsutsumi, S., Ohno, K., Yamada, J., Kouno, E., Kumata, H., Distribution of polycyclic aromatic hydrocarbons (PAHs) in rivers and estuaries in Malaysia - A widespread input of petrogenic PAHs (2002) Environ. Sci. Technol, 36, pp. 1907-1918
- Windsor Jr, J.H., Hites, R.A., Polycyclic aromatic hydrocarbons in Gulf of Maine sediment and Nova Scotia soils (1979) Geochim. Cosmochim. Acta, 43, pp. 27-33

- Shiaris, M.P., Jambard-Sweet, D., Distribution of polycyclic aromatic hydrocarbons in surficial sediments of Boston Harbor, Massachusetts, USA (1986) Mar. Pollut. Bull, 17, pp. 469-472
- Van Metre, P.C., Mahler, B.J., Furlong, E.T., Urban sprawl leaves its PAH signature (2000) Environ. Sci. Technol, 34, pp. 4064-4070
- Phuong, P.K., Son, C.P.N., Sauvain, J.J., Tarradellas, J., Contamination of PCB's, DDT's, and heavy metals in sediment of Ho Chi Minh City's canals, Vietnam (1998) Bull. Environ. Contam. Toxicol, 60, pp. 347-354
- (1991) Manual for monitoring water and sediment in the environment, , Ministry of the Environment, Government of Japan , in Japanese
- Baxtor, R.M., Reductive dechlorination of certain chlorinated organic compounds by reduced hematin compared with their behavior in the environment (1990) Chemosphere, 121, pp. 451-458
- DDT and its derivatives-environmental aspects (1989) Environmental Health Criteria 83, , World Health Organization , Geneva
- Li, Y.F., Cai, D.J., Singh, A., Technical hexachlorocyclohexane use and trend in China and their impact on the environment (1998) Arch. Environ. Contam. Toxicol, 35, pp. 688-697
- Fu, S., Chu, S., Xu, X., Organochlorine pesticide residue in soils from Tibet, China (2001) Bull. Environ. Contam. Toxicol, 66, pp. 171-177
- Hung, D.Q., Thiemann, W., Contamination by selected chlorinated pesticides in surface waters in Hanoi. Vietnam (2002) Chemosphere, 47, pp. 357-367
- Patton, G.W., Hinckley, D.A., Walla, M.D., Bidleman, T.F., Airborne organochlorines in the Canadian High Arctic (1989) Tellus, 41 B, pp. 243-255
- Hoff, R.M., Chan, K.W., Atmospheric concentrations of chlordane at Mould Bay, N.W.T., Canada (1986) Chemosphere, 15, pp. 449-452
- Takasuga, T., Inoue, T., Ohi, E., All congener specific analytical method for polychlorinated biphenyls with various chromatographic clean-up and HRGC/HRMS (1995) J. Environ. Chem, 5, pp. 647-675. , in Japanese
- Alder, A.C., Haggblom, M.M., Oppenhelmer, S.R., Yong, L.Y., Reductive dechlorination of polychlorinated biphenyls in anaerobic sediments (1993) Environ. Sci. Technol, 27, pp. 530-538
- Sokol, R.C., Bethoney, C.M., Rhee, G.-Y., Effect of PCB concentration on reductive dechlorination and dechlorination potential in natural sediment (1995) Water Res, 29, pp. 45-48
- Hartkamp-Commandeur, L.C.M., Gerritse, J., Govers, H.A.J., Parsons, J.R., Reductive dehalogenation of polychlorinated biphenyls by anaerobic microorganisms enriched from Dutch sediments (1996) Chemosphere, 32, pp. 1275-1286
- Vanier, C., Sylvestre, M., Planas, D., Persistence and fate of PCBs in sediment of the Saint Lawrence River (1996) Sci. Total Environ, 102, pp. 229-244
- Hites, R.A., Laflamme, R.E., Windsor, J.G., Farrington, J.W., Deuser, W.D., Polycyclic aromatic hydrocarbons in an anoxic sediment core from the Pettaquamscutt River (Rhode Island, USA) (1980) Geochim. Cosmochim. Acta, 44, pp. 873-878